

BATES (W.H.)


ON THE NECESSITY OF
SUFFICIENTLY EXTENDED
URETHRAL INCISIONS

FOR THE
PERMANENT CURE OF STRICTURES OF
THE URETHRA.

*WITH A DESCRIPTION OF A NEW URETHROTOME OF
VERY SMALL CALIBRE, TO BE USED PREPARATORY
TO DR. OTIS'S DILATING URETHROTOME.*

BY
W. H. BATES, M. D.,
BROOKLYN.

[REPRINTED FROM THE NEW YORK MEDICAL JOURNAL, JAN., 1877.]



NEW YORK:
D. APPLETON & COMPANY,
549 & 551 BROADWAY.
1877.

APPLETONS' PERIODICALS.

APPLETONS' JOURNAL:

A MONTHLY MISCELLANY OF POPULAR LITERATURE.

NEW SERIES.

TWENTY-FIVE CENTS PER NUMBER. THREE DOLLARS PER ANNUM.

APPLETONS' JOURNAL is now published monthly; it is devoted to popular literature and all matters of taste and general culture—published at a price to bring it within the reach of all classes. It contains superior fiction, in the form of serials and short stories; papers graphically descriptive of picturesque places; articles upon men of note, and upon the habits of different peoples; essays upon household and social topics; articles of travel and adventure; scientific and industrial articles written in a graphic and popular style. In brief, the aim is to be comprehensive, including in its plan all branches of literature and all themes of interest to intelligent readers. Each number is illustrated.

TERMS: Three dollars per annum, postage prepaid, to all subscribers in the United States; or Twenty-five Cents per number. A Club of Four Yearly Subscriptions will entitle the sender to an extra subscription gratis; that is, five copies will be sent one year for twelve dollars. For \$7.20, APPLETONS' JOURNAL and THE POPULAR SCIENCE MONTHLY (full price, eight dollars), postage prepaid.

THE POPULAR SCIENCE MONTHLY.

Conducted by E. L. YOUMANS.

This periodical was started (in 1872) to promote the diffusion of valuable scientific knowledge, in a readable and attractive form, among all classes of the community, and has thus far met a want supplied by no other magazine in the United States.

Containing instructive and interesting articles and abstracts of articles, original, selected, translated, and illustrated, from the pens of the leading scientific men of different countries; accounts of important scientific discoveries, the application of science to the practical arts, and the latest views put forth concerning natural phenomena, have been given by *savants* of the highest authority. Prominent attention has been also devoted to those various sciences which help to a better understanding of the nature of man, to the bearings of science upon the questions of society and government, to scientific education, and to the conflicts which spring from the progressive nature of scientific knowledge.

THE POPULAR SCIENCE MONTHLY is published monthly in a large octavo, handsomely printed on clear type, and, when the subjects admit, fully illustrated.

TERMS: \$5.00 per annum, or Fifty Cents per Number. Postage prepaid to all Subscribers in the United States.

THE ART JOURNAL:

An International Gallery of Engravings,

BY DISTINGUISHED ARTISTS OF EUROPE AND AMERICA;

WITH ILLUSTRATED PAPERS IN THE VARIOUS BRANCHES OF ART.

THE ART JOURNAL is a monthly publication, quarto size, superbly illustrated and printed, and specially devoted to the world of Art—Painting, Sculpture, Architecture, Decoration, Engraving, Etching, Enameling, and Designing in all its branches—having in view the double purpose of supplying a complete illustrated record of progress in the Arts, and of affording a means for the cultivation of Art-taste among the people. Each number is richly and abundantly illustrated on both steel and wood, and no pains are spared to render this "ART JOURNAL" the most valuable publication of the kind in the world. It contains the Steel Plates and Illustrations of the LONDON ART JOURNAL, a publication of world-wide fame (the exclusive right of which, for Canada and the United States, has been purchased by the publishers); with *exclusive additions devoted principally to American Art and American topics. Published monthly. Sold only by Subscription.* Price, 75 Cents per Number; \$9.00 per Annum, postage prepaid.

Subscriptions received by the Publishers, or their Agents. AGENCIES: 22 Hawley St., Boston; 922 Chestnut St., Philadelphia; 22 Post-Office Avenue, Baltimore; 53 Ninth St., Pittsburg; 100 State St., Albany; 42 State St., Rochester; 103 State St., Chicago; 30 W. 4th St., Cincinnati; 305 Locust St., St. Louis; 20 St. Charles St., New Orleans; 230 Sutter St., San Francisco.

D. APPLETON & CO., Publishers, 549 & 551 Broadway, N. Y.


ON THE NECESSITY OF
SUFFICIENTLY EXTENDED
URETHRAL INCISIONS

FOR THE
PERMANENT CURE OF STRICTURES OF
THE URETHRA.

*WITH A DESCRIPTION OF A NEW URETHROTOME OF
VERY SMALL CALIBRE, TO BE USED PREPARATORY
TO DR. OTIS'S DILATING URETHROTOME.*

BY
W. H. BATES, M. D.,
BROOKLYN.

[REPRINTED FROM THE NEW YORK MEDICAL JOURNAL, JAN., 1877.]



NEW YORK:
D. APPLETON & COMPANY,
549 & 551 BROADWAY.
1877.

ON THE NECESSITY OF SUFFICIENTLY
EXTENDED URETHRAL INCISIONS FOR
THE PERMANENT CURE OF STRICTURES
OF THE URETHRA. WITH A DESCRIP-
TION OF A NEW URETHROTOME OF
VERY SMALL CALIBRE, TO BE USED
PREPARATORY TO DR. OTIS'S DILATING
URETHROTOME.

PROBABLY at no time has the subject of the treatment of strictures of the urethra attracted more attention from surgeons than the present—more especially since the revival of the operation of internal urethrotomy, and the introduction of new instruments for the purpose of diagnosis and treatment of this class of troublesome affections—not only to the patients, but to the surgeon. It must be obvious that the value of all new principles advanced, and the curative procedures under them, cannot be judged or criticised from an abstract point of view; therefore practical experience upon real cases is the only reliable basis upon which rational conclusions can be founded. With this view I propose to contribute some of the results of my personal experience, which bear upon the recent discussions in this department of sur-

gery—excluding such details as are general to strictures as a class.

The most protracted case I have any record of is that of a man now over seventy-four years of age, on whom I operated between eight and nine years ago. At that time his stricture had existed for nearly fifty years. He had been under the treatment of some of the ablest surgeons, including Drs. Valentine Mott and J. Kearney Rodgers. He had been in the habit of passing a No. 16 French steel sound almost daily. After several months' treatment, I did not succeed in carrying dilatation above No. 18 French, which was painful, and as a last resort I divided the stricture with Civiale's urethrotome, after which No. 27 French steel sound was readily introduced into his bladder. The introduction of this same number was continued for some time with marked benefit. The patient's general condition being greatly improved, his urethral discharge diminished. I then lost sight of him for a time; subsequently I saw him for other trouble; he informed me that he was very much better. Last spring he came to my office, and I desired to examine his urethra. He stated that he had remained well. I passed easily the same sound I had used on him previously. I also examined him with the urethrometer, and found no trace of stricture at the seat of division. The patient had found no necessity for having recourse to any instruments, since those passed by myself.

The next case was likewise a protracted one. Mr. K., aged thirty-three years. For five years past has been troubled with muco-purulent discharge, which would occasionally cease for a month or two, and then return, without apparent cause. Had been constantly under treatment, and examined for stricture by several surgeons, but was assured that none existed. On examination, I found a stricture of large calibre, readily admitting a No. 27 steel sound, about four inches from the meatus. Persevering efforts to effect higher dilatation were continued through several months, with only temporary benefit, the stricture remaining firm and rigid. Divulsion by means of Holt's divulsor was resorted to, but without permanent benefit. I then divided the stricture with a urethrotome, up

to No. 30 French, and introduced No. 30 steel sound into the bladder daily, for a few days, when the patient informed me that all discharge had ceased, and he discontinued his attendance. He has remained perfectly well up to the present time, now over two years, using no sound whatever. I quite recently examined his urethra carefully with the urethrometer and bulbous sounds, and failed to detect the slightest indication of stricture.

The next case was one of multiple stricture. Mr. R., aged twenty-five years. Six or seven years ago he contracted gonorrhœa, which ran into a gleet, for which he was constantly treated; came under my care more than a year ago. Three strictures were detected and operated upon successively, with Dr. Otis's dilating urethrotome, dividing little above the normal calibre, thirty-four millimetres, previously ascertained with the urethrometer. The rapid cessation of the discharge followed the operations. This patient has frequently been carefully examined since, and there is no evidence of any stricture left. He has likewise had no subsequent recourse to instruments.

The next case illustrates the persistence of stricture with free discharge, notwithstanding what would be considered an ample urethra. (*Vide* Sir Henry Thompson's recent lectures, London *Lancet*.) Mr. M., aged thirty-five years. Had been troubled with stricture for several years, for which he had been treated by a surgeon in Germany, by dilatation, which he had very irregularly continued. Recontraction had taken place, followed by free discharge, for which I was consulted. After the acute symptoms had subsided, the normal calibre of the urethra, thirty-seven millimetres, was restored by operation, the strictured portion being thirty millimetres. This patient has remained well ever since, now over a year. Uses no sound.

The following case is of interest, showing the effect of the restoration of the urethra on long-continued discharge, the discharge in this case having lasted for a period of twenty-six years. Mr. F., aged forty-five years. Contracted gonorrhœa at the age of nineteen; over a year ago was placed under my

care for treatment by Dr. Shaw. Stated that he had never been free from urethral discharge since he first contracted disease. On examination, four strictures were found, three of which, situated in the pendulous portion, were divided, and the normal calibre of the urethra restored. The fourth, situated in the membranous portion, was divulsed. The frequent introduction of sounds for a few weeks was continued. This patient was so much pleased with the relief given him (having frequently suffered from retention), that he declined to submit to anything being done for the cessation of the discharge, considering it of no consequence. A short time ago he called upon me, and I examined him, and found that not the slightest contraction had taken place in the strictures that had been divided. One situated in the membranous portion which had been divulsed, but not cut, had recontracted (of which I shall speak hereafter). There was not the slightest trace of any discharge to be detected from this patient's urethra. He informed me that it had recently entirely ceased. There had been no introduction of instruments in this case after the first few weeks subsequent to the operation. I think this case illustrates somewhat the effects of the operation upon a similar case of long-continued discharge, which Dr. Otis operated on in July, in London, for the cure of gleet, and which Mr. Berkeley Hill published in the *Lancet*. He stated that "the discharge ran on until the holidays, and was then cured by other means." In that case, as in this, the subsequent cessation of the discharge may have been due to the original operation, and not to the after-medication.

The most interesting feature of the next case which I shall offer is the very large calibre of the urethra, coexisting with the persisting symptoms of stricture. I wish it understood that these figures are not the result of dilatation, but simply the normal measurement, as determined by the urethrometer and bulbous sounds. Mr. W., aged thirty-three years; height five feet eleven inches, weight one hundred and eighty pounds. This patient contracted a gonorrhœa two or three years ago, followed by a gleet. The discharge never entirely ceased. On examination, found a stricture at the meatus, of twenty-five

millimetres calibre. The normal calibre of the urethra, as ascertained by the urethrometer, was forty-five millimetres. A second stricture was found four inches from the meatus, of a calibre of thirty-seven millimetres. The strictures were divided so as freely to admit No. 40 steel sound, being the largest then in my possession, which was passed daily, until the parts healed.

The discharge rapidly diminished. The patient remained under treatment for several weeks, but the discharge not disappearing, although several operations had been performed with Otis's dilating urethrotome expanded to its fullest extent, forty-six millimetres, and a large bulbous bougie passed every second day to keep the incision open, without much benefit, the patient, being desirous of returning to his home in Massachusetts, promised to correspond with me, and, if no improvement took place, to return again at some future time for further treatment.

He wrote me a couple of times, saying that he remained about the same, with the exception of the stricture at four inches, which failed to admit his bulbous bougie so readily as formerly (which I had, contrary to my usual habit, requested him to pass occasionally).

He did return to the city early in the autumn for further treatment.

I operated on him again for the second stricture, which had recontracted, but still without satisfactory results.

I now determined to divide the stricture-tissue more deeply, feeling satisfied that my incisions had failed to completely sever the stricture, owing to the largest urethrotome being too small to divide it completely. After a little time, I devised the following means of increasing the size of Otis's dilating urethrotome: I took a piece of wood cut to the size and shape of the lower blade of the instrument, and with rubber tubing one-eighth of an inch in diameter, binding the wood firmly, and completely covering it. This gave me a perfectly smooth and even surface, and increased the size of the urethrotome seven millimetres by measurement.

With this addition I operated, dilating and dividing up to

fifty-three millimetres. There was very little hæmorrhage following this deep incision—certainly not more than two drachms. The patient had provided himself with a steel sound of forty-five millimetres calibre, which was introduced daily for about a week. The patient rapidly improved after this last operation, the discharge ceasing entirely. He has returned home, promising to communicate with me again if there should be any recurrence. On careful examination with bulbous sounds, I failed to detect the slightest contraction. Of course, it is too soon to speak as to the ultimate result in this case. I report it at this time simply on account of the very large size of the urethral canal, admitting No. 45 steel sound to pass readily into the bladder, by simply guiding it. *Should* such a thing recur again, such is my faith in the principles here enunciated and illustrated that, notwithstanding the dilatation hitherto attained, I should dilate and divide until all resistance ceased, and morbid symptoms disappeared; after this has been accomplished, it will be quite time enough to use medication, which, it is my growing conviction, by these procedures will be rendered unnecessary.

The cases narrated by no means exhaust the records which have accumulated in my possession, but I think at this time it is unnecessary to add more, as those that I have selected are not exceptional, but the average results I have obtained by internal urethrotomy. I am satisfied, from my own experience, that the operation fails sometimes, owing to its incomplete performance.

As to the future of these cases, that still remains in doubt; and, if hereafter I find that recontraction has taken place, I shall be pleased to report such fact.

To none of my patients on whom I operate do I advise the continued use of sounds; with what results, the few examples given may serve as illustration.

In common with all surgeons, I have occasionally met with strictures of small calibre, that it was desirable to divide, in order to allow the introduction of a sufficiently large urethrotome, to complete the division, for the purpose of effecting

a cure; and have felt the want of a sufficiently small and safe urethrotome suitable in these cases.

The best that we had was Maisonneuve's, which cuts from before backward, and is liable to get out of the urethra, thus making a troublesome and often dangerous false passage. To avoid this, Dr. Gouley has added a "bridged" tip, so that it threads over a whalebone guide, but the blade is not concealed, and is liable to wound the urethra where it should not. M. Horteloup, of Paris, has very recently devised a new urethrotome, or set of urethrotomes, comprising four instruments (*see Annales de Dermatologie et de Syphiligraphie*, tome vii., page 453), which have concealed blades—the smallest of which cuts to eleven millimetres, and the largest to twenty-three millimetres. There is a bulb projecting on the under surface of these instruments, where the blade is concealed. This is claimed to be a great improvement on Maisonneuve's but, like it, cuts from before backward, which is objectionable; another objection to this novelty is, that four separate instruments are required to effect the object, which is quite as thoroughly attained by the single one, which I now propose to describe. Let me premise that this instrument was manufactured from my plans, and under my direction, by Messrs. Shepard & Dudley, of New York.

The drawings for it were made over a year ago, and the instrument has been in my hands for a long time. I have waited for suitable cases on which to test it previous to making it public, in order to remedy any existing defects that might be disclosed in its use. I offer it simply as an adjunct to Dr. Otis's dilating urethrotome, believing that it is better and safer than any of the existing small urethrotomes in cases requiring its use. It consists of a handle to which is attached a steel canula, eight inches long, containing a concealed knife, which may be projected to the extent of twenty millimetres. The instrument terminates in a "bridged" tip, which screws on the canula; there are two of these tips accompanying each instrument, pointing in opposite directions for the purpose of enabling the operator to use the knife either on the upper or under surface of the urethra, as he may deem advisable, this still being a question *sub judice*.

The blade is concealed in the end of the canula, and is projected by turning the screw in the end of the handle to the right. The end of the canula where the knife is projected through the slot is of solid metal, rendering firm support to the blade, and adding greatly to the strength of the instrument. To the handle is attached a scale, with an index, such as is used by Dr. Otis on his instruments, which enables the operator to know and govern the exact limits of his incision. The diameter of this instrument at its tip is two and two-thirds millimetres. The canula is a little more, but considerably less than three millimetres, bringing the instrument down to about the calibre of Maisonneuve's. It is operated in this manner: After having passed one of Otis's long whalebone guides through the stricture into the bladder, the tip of the instrument is threaded over it, and passed through the stricture. The knife is then projected by turning the screw in the handle to the right. The instrument is then slowly withdrawn; as soon as all resistance ceases, the blade is withdrawn into the canula, by reversing the screw in the handle, viz., to the left.

This will prevent the possibility of wounding the healthy portion of the urethra; in other words, the knife cuts only where the operator wills it.

The advantages of this instrument are, that it cuts from behind forward; consequently there is no danger of getting out of the urethra. It combines simplicity with strength, is easily cleansed and kept in order, and last, but not least, is not more costly than any urethrotome of small calibre hitherto offered to the profession.

An examination of the accompanying woodcut, which represents the knife half closed, will render more minute descriptive details unnecessary.

Its special applicability was most manifest in the case of Mr. F., already described. It will be remembered that



this patient had a stricture in the membranous portion of the urethra, which had been ineffectually divulsed, while three situated anterior to it had been successfully cut without subsequent contraction. It is a well-known fact that the division of strictures in the deep urethra is attended with greater danger from hæmorrhage than those made in other parts more anterior. Hence the importance of confining the forme within strict limits of morbid structure. This instrument enables the operator so to appreciate the parts to be cut that all divisions necessary to be made reduce the danger to a minimum. Thus, in Mr. F.'s case, when the stricture already divulsed had recontracted until a small whalebone guide could only be passed through it, upon this the urethrotome was threaded, and the stricture divided to twenty millimetres, after which Otis's dilating urethrotome was readily introduced, and the canal enlarged to thirty-two millimetres. So little hæmorrhage followed that no artificial means were required for its arrest, and, after thirty-six hours confinement in bed, he returned to his accustomed business. Since then the patient has remained perfectly well, and free from inconvenience.

MEDICAL WORKS

PUBLISHED BY D. APPLETON & CO.

<i>Anstie on Neuralgia.</i> 1 vol., 12mo.....	Cloth, \$2 50
<i>Bartholow's Treatise on Therapeutics</i>	" 5 00
<i>Barker on Puerperal Diseases.</i> 1 vol.....	" 5 00
<i>Barker on Sea-Sickness.</i> 1 vol., 16mo.....	" 75
<i>Barnes's Obstetric Operations.</i> 1 vol., 8vo.....	" 4 50
<i>Bellevue and Charity Hospital Reports.</i> 1 vol., 8vo.....	" 4 00
<i>Bennet's Winter and Spring on the Mediterranean.</i> 1 vol., 12mo.....	" 3 50
<i>Bennet on the Treatment of Pulmonary Consumption.</i> 1 vol., 8vo.....	" 1 50
<i>Billroth's General Surgical Pathology and Therapeutics.</i> 1 vol., 8vo.....	" 5 00
<i>Buck's Contributions to Reparative Surgery.</i> 1 vol., 8vo.....	" 3 00
<i>Bastian on the Common Forms of Paralysis from Brain Diseases</i>	" 1 75
<i>Bulkeley's (L. D.) Acne; its Pathology, etc.</i>	(In press.)
<i>Combe on the Management of Infancy.</i> 1 vol., 12mo.....	Cloth, 1 50
<i>Carpenter's Mental Physiology</i>	" 3 00
<i>Chauveau's Comparative Anatomy of the Domesticated Animals.</i> Edited by George Fleming, F. R. G. S., M. A. I. 1 vol., 8vo, with 450 Illustrations.....	" 6 00
<i>Davis's (Henry G.) Conservative Surgery</i>	" 3 00
<i>Dielskon on Medicine in Relation to the Mind</i>	" 3 50
<i>Elliot's Obstetric Clinic.</i> 1 vol., 8vo.....	" 4 50
<i>Echcr's Convolutions of the Brain</i>	" 1 25
<i>Flint's Physiology.</i> 5 vols., 8vo.....	Cloth, per vol., \$4 50; Sheep, 5 50
<i>Flint's Text-Book of Human Physiology.</i> 1 vol., 8vo.....	Cloth, \$6 00; " 7 00
<i>Flint's Manual on Urine.</i> 1 vol., 12mo.....	Cloth, 1 00
<i>Flint's Relations of Urea to Exercise.</i> 1 vol., 8vo.....	" 1 00
<i>Frey's Histology and Histo-Chemistry of Man</i>	" 5 00
<i>Hoffmann's Manual of Medicinal Chemicals</i>	" 3 00
<i>Holland's (Sir Henry) Recollections of Past Life.</i> 1 vol., 12mo.....	" 2 00
<i>Howe on Emergencies.</i> 1 vol., 8vo.....	" 3 00
<i>Howe on the Breath, and the Diseases which give it a Fetid Odor</i>	" 1 00
<i>Huxley on the Anatomy of Vertebrated Animals.</i> 1 vol.....	" 2 50
<i>Huxley and Youmans's Physiology and Hygiene.</i> 1 vol., 12mo.....	" 1 75
<i>Hammond's Insanity in its Relations to Crine.</i> 1 vol., 8vo.....	" 1 00
<i>Hammond's Diseases of the Nervous System.</i> 1 vol., 8vo.....	Cloth, \$6 00; Sheep, 7 00
<i>Hammond's Clinical Lectures on Diseases of the Nervous System.</i> 1 vol., 8vo.....	" 3 50
<i>Hamilton's (A. McL.) Electro-Therapeutics.</i> 1 vol., 8vo.....	Cloth, 2 00
<i>Johnston's Chemistry of Common Life.</i> 2 vols., 12mo.....	" 3 00
<i>Letterman's Recollections of the Army of the Potomac.</i> 1 vol., 8vo.....	" 1 00
<i>Lewes's Physiology of Common Life.</i> 2 vols., 12mo.....	" 3 00
<i>Markoe on Diseases of the Bones.</i> 1 vol., 8vo.....	" 4 50
<i>Maudsley on the Mind.</i> 1 vol., 8vo.....	" 3 50
<i>Maudsley's Body and Mind.</i> 1 vol., 12mo.....	" 1 00
<i>Maudsley on Responsibility in Mental Disease</i>	" 1 50
<i>Meyer's Electricity.</i> 1 vol., 8vo.....	" 4 50
<i>Niemeyer's Practical Medicine.</i> 2 vols., 8vo.....	Cloth, \$9 00; Sheep, 11 00
<i>Nefel on Galvano-Therapeutics.</i> 1 vol., 12mo.....	Cloth, 1 50
<i>Nightingale's Notes on Nursing.</i> 1 vol., 12mo.....	" 75
<i>Neumann on Skin Diseases.</i> 1 vol., 8vo.....	" 4 00
<i>New York Medical Journal</i>	\$4 00 per annum. Specimen copies, 35
<i>Paget's Clinical Lectures and Essays.</i> 1 vol., 8vo.....	Cloth, 5 00
<i>Parslee on Ovarian Tumors.</i> 1 vol., 8vo.....	" 5 00
<i>Pereira's Materia Medica and Therapeutics.</i> 1 vol., 8vo.....	Cloth, \$7 00; Sheep, 8 00
<i>Richardson's Diseases of Modern Life.</i> 1 vol., 12mo.....	Cloth, 2 00
<i>Sayre's Club-Foot.</i> 1 vol., 12mo.....	" 1 00
<i>Sayre's Orthopedic Surgery.</i> With the Operations incident to Deformities. With numerous Illustrations.....	Cloth, \$5 00; Sheep, 6 00
<i>Schroeder on Obstetrics.</i> 1 vol., 8vo.....	Cloth, 3 50
<i>Steiner's Compendium of Children's Diseases</i>	" 3 50
<i>Stroud's Physical Cause of the Death of Christ.</i> 1 vol., 12mo.....	" 2 00
<i>Swett on Diseases of the Chest.</i> 1 vol., 8vo.....	" 3 50
<i>Simpson's (Sir Jas. Y.) Complete Works.</i> Vol. I. Obstetrics and Gynecology. 8vo. Vol. II. Anesthesia, Hospitalism, etc. 8vo. Vol. III. The Diseases of Women.....	Per vol. Cloth, \$3 00; Sheep, 4 00
<i>Tilt's Uterine Therapeutics.</i> 1 vol., 8vo.....	Cloth, 3 50
<i>Van Buren on Diseases of the Rectum.</i> 1 vol., 12mo.....	" 1 50
<i>Van Buren & Keyes's Genito-Urinary Diseases, with Syphilis.</i> Cloth, \$5; Sheep, 6 00	
<i>Vogel's Diseases of Children.</i> 1 vol., 8vo.....	Cloth, \$4 50; " 5 50
<i>Wells on Diseases of the Ovaries.</i> 1 vol., 8vo.....	Cloth, 5 00
<i>Wagner's Chemical Technology.</i> 1 vol., 8vo.....	" 5 00
<i>Walton's Mineral Springs of the United States and Canada.</i> With Analyses and Notes on the Prominent Spas of Europe.....	" 2 00

* * Any of these works will be mailed, post-free, to any part of the United States, on receipt of the price.

A large and carefully-selected stock of Medical Works, American and Foreign, constantly on hand. Descriptive Catalogue forwarded on application.

Physicians desiring to have their names inserted in our Medical Directory of the United States and Canada, will please send them in full, with addresses. No charge.

D. APPLETON & CO., Publishers, 549 & 551 Broadway, New York.

APPLETONS', AMERICAN CYCLOPÆDIA.

NEW REVISED EDITION.

*Entirely rewritten by the ablest writers on every subject. Printed from new type,
and illustrated with Several Thousand Engravings and Maps.*

The work originally published under the title of *THE NEW AMERICAN CYCLOPÆDIA* was completed in 1863, since which time the wide circulation which it has attained in all parts of the United States, and the signal developments which have taken place in every branch of science, literature, and art, have induced the editors and publishers to submit it to an exact and thorough revision, and to issue a new edition entitled *THE AMERICAN CYCLOPÆDIA*.

Within the last ten years the progress of discovery in every department of knowledge has made a new work of reference an imperative want.

The movement of political affairs has kept pace with the discoveries of science, and their fruitful application to the industrial and useful arts and the convenience and refinement of social life. Great wars and consequent revolutions have occurred, involving national changes of peculiar moment. The civil war of our own country, which was at its height when the last volume of the old work appeared, has happily been ended, and a new course of commercial and industrial activity has been commenced.

Large accessions to our geographical knowledge have been made by the indefatigable explorers of Africa.

The great political revolutions of the last decade, with the natural result of the lapse of time, have brought into public view a multitude of new men, whose names are in every one's mouth, and of whose lives every one is curious to know the particulars. Great battles have been fought, and important sieges maintained, of which the details are as yet preserved only in the newspapers, or in the transient publications of the day, but which ought now to take their place in permanent and authentic history.

In preparing the present edition for the press, it has accordingly been the aim of the editors to bring down the information to the latest possible dates, and to furnish an accurate account of the most recent discoveries in science, of every fresh production in literature, and the newest inventions in the practical arts, as well as to give a succinct and original record of the progress of political and historical events.

The work has been begun after long and careful preliminary labor, and with the most ample resources for carrying it on to a successful termination.

None of the original stereotype plates have been used, but every page has been printed on new type, forming in fact a new Cyclopædia, with the same plan and compass as its predecessor, but with a far greater pecuniary expenditure, and with such improvements in its composition as have been suggested by longer experience and enlarged knowledge.

The illustrations, which are introduced for the first time in the present edition, have been added not for the sake of pictorial effect, but to give greater lucidity and force to the explanations in the text. They embrace all branches of science and of natural history, and depict the most famous and remarkable features of scenery, architecture, and art, as well as the various processes of mechanics and manufactures. Although intended for instruction rather than embellishment, no pains have been spared to insure their artistic excellence; the cost of their execution is enormous, and it is believed that they will find a welcome reception as an admirable feature of the Cyclopædia, and worthy of its high character.

This work is sold to subscribers only, payable on delivery of each volume. It is now completed in sixteen large octavo volumes, each containing over 800 pages, fully illustrated with several thousand Wood Engravings, and with numerous colored Lithographic Maps.

PRICE AND STYLE OF BINDING.

<i>In extra cloth, per vol.</i> \$5.00	<i>In half russia, extra gilt, per vol.</i> . . \$8.00
<i>In library leather, per vol.</i> 6.00	<i>In full morocco antique, gilt edges, per vol.</i> 10.00
<i>In half turkey morocco, per vol.</i> 7.00	<i>In full russia, per vol.</i> 10.00

* * * Specimen pages of the *AMERICAN CYCLOPÆDIA*, showing type, illustrations, etc., will be sent gratis, on application.

D. APPLETON & CO., PUBLISHERS,

549 & 551 Broadway, New York.